

Adjoint Semilattice and Minimal Brouwerian Extensions of a Hilbert Algebra*

Jānis CĪRULIS

Faculty of Computing, University of Latvia,
Raina b. 19, Riga LV-1586, Latvia
e-mail: jc@lanet.lv

(Received October 30, 2011)

Abstract

Let $A := (A, \rightarrow, 1)$ be a Hilbert algebra. The monoid of all unary operations on A generated by operations $\alpha_p: x \mapsto (p \rightarrow x)$, which is actually an upper semilattice w.r.t. the pointwise ordering, is called the adjoint semilattice of A . This semilattice is isomorphic to the semilattice of finitely generated filters of A , it is subtractive (i.e., dually implicative), and its ideal lattice is isomorphic to the filter lattice of A . Moreover, the order dual of the adjoint semilattice is a minimal Brouwerian extension of A , and the embedding of A into this extension preserves all existing joins and certain “compatible” meets.

Key words: adjoint semilattice, Brouwerian extension, closure endomorphism, compatible meet, filter, Hilbert algebra, implicative semilattice, subtraction

2000 Mathematics Subject Classification: 03G25, 06A12, 06A15, 08A35

1 Introduction

Let $A := (A, \rightarrow, 1)$ be a Hilbert algebra. A mapping $\varphi: A \rightarrow A$ is called a *closure endomorphism* if it is simultaneously a closure operator and an endomorphism.

This notion goes back to [15], where Glivenko operators on implicative, or Brouwerian, semilattices were discussed. In [16], it was shown that the Glivenko operators are precisely the closure endomorphisms and that all such endomorphisms form a distributive lattice. Connections of this lattice with the filter

* Supported by ESF Project 2009/0216/1DP/1.1.1.2.0/09/APIA/VIAA/044

lattice and with a certain sublattice of subalgebras of an implicative semilattice were discovered in [17].

Closure endomorphisms on Hilbert algebras were introduced by the present author in [1] and further studied in [2]. The identity mapping ε , the unit mapping $\iota: x \mapsto 1$ and, for any $p \in A$, the mappings α_p and β_p defined by

$$\alpha_p x := p \rightarrow x, \quad \beta_p x := (x \rightarrow p) \rightarrow x,$$

respectively are examples of closure endomorphisms. The set CE of all closure endomorphisms on A is closed under composition \circ and pointwise defined meets. The algebra $(CE, \circ, \wedge, \varepsilon, \iota)$ also is a bounded distributive lattice [2], in which \circ acts as join and the natural ordering may be defined pointwise. Furthermore, an endomorphism φ is a closure operator if and only if

$$\varphi(x \rightarrow y) = x \rightarrow \varphi y. \quad (1)$$

In this paper, we pay attention to closure endomorphisms α_p and their compositions. For every finite subset $P := \{p_1, p_2, \dots, p_n\}$ of A (in symbols, $P \subseteq_{\text{fin}} A$), we set $\alpha_P := \alpha_{p_n} \circ \dots \circ \alpha_{p_2} \circ \alpha_{p_1}$ (we shall usually drop the symbol ‘ \circ ’ in notation). In the case when P is empty, this means that $\alpha_P = \varepsilon$. Of course, each mapping α_P also is a closure endomorphism; we shall call them *finitely generated* (cf. Proposition 2 below). The set CE^f of all such mappings is closed under composition, and the algebra $(CE^f, \circ, \varepsilon)$ is a lower bounded join semilattice. In the dual context of BCI/BCK-algebras, the counterpart of CE^f is usually called the adjoint semigroup (or monoid) of an algebra under consideration (see, for example, [8, 9, 13]). We adopt this term and call CE^f the *adjoint semilattice* of the initial Hilbert algebra A . It is shown in Section 3 to be isomorphic to the semilattice of finitely generated filters of A and subtractive, i.e., dually implicative, while its generating set turns out to be closed under subtraction and is an order dual of A (Section 4). The lattice of ideals of CE^f is isomorphic to the lattice of filters of A (Section 3). A minimal Brouwerian extension of A is a minimal implicative semilattice of which A is a subreduct; in Section 4 such an extension is shown to be dually isomorphic to the adjoint semilattice of A . Embedding of A into its minimal Brouwerian extension preserves all existing joins; we characterize also the preserved meets (Section 5).

2 Preliminaries

We assume that the reader is acquainted with the notion of Hilbert algebra and with elementary arithmetics in such algebras. This information can be found, e.g., in [2, 3, 5, 6, 14]. Recall that Hilbert algebras were introduced in [5] as the order duals of L. Henkin’s *implicational models* [6]. In [2, 3] also the notion of compatible meet (suggested by [14]) in a Hilbert algebra was introduced and discussed. We now list some basic facts concerning it.

Let $A := (A, \rightarrow, 1)$ be a Hilbert algebra. Elements $a, b \in A$ are said to be *compatible* (in symbols, $a \mathcal{C} b$) if there is a lower bound c of $\{a, b\}$ such that

$a \leq b \rightarrow c$. If this is the case, then c is the g.l.b. of a and b ; we call this element the *compatible meet* of a and b and denote it by $a \wedge b$. In this way, we come to a partial operation \wedge on A . It is total if and only if A is actually an implicative semilattice. For example, if $\varphi, \psi \in CE$, then always $\varphi x C \psi x$. A *relative subsemilattice* of A is any subset of A closed under existing compatible meets. Thus, the subset $CE(a) := \{\varphi a : \varphi \in CE\}$ is a relative subsemilattice for every $a \in A$: all meets in it exist and are compatible. (In [2, 3], we used the notation xy for the meet of x and y , and wrote $x \wedge y$ for it if it was compatible.)

Examples of relative subsemilattices are provided also by filters. A *filter* (an implicative filter, a deductive system) of A is a subset J containing 1 and such that $y \in J$ whenever $x, x \rightarrow y \in J$. According to [3, Lemma 3.2], J is an implicative filter if and only if it is a semilattice filter, i.e., an upwards closed relative subsemilattice of A .

The above definition of compatibility is equivalent to the original one presented in [14]. It was also observed there that elements x and y are compatible if and only if the filter generated by $\{x, y\}$ is principal, and then $x \wedge y$ is the least element of the filter. We now consider an arbitrary finite subset $P \subseteq A$ as *compatible* if P has a lower bound c (necessary unique) in $[P]$, and say that then c is the *compatible meet of P* (denoted by $\bigwedge P$). This is the case if and only if $[P] = [c]$. Equivalently, a subset P is compatible if $\alpha_P = \alpha_p$ for some $p \in A$. In particular, the empty set is compatible; of course, $\bigwedge \emptyset = 1$.

Where $X \subseteq A$, we denote by $\alpha_P(X)$ the set $\{\alpha_P x : x \in X\}$. A one-element subset of A is identified with its single element. The following notation will be convenient (cf. Definition 6.4 in [7]): given two finite subsets, P and Q , of A , we shall write $Q \rightarrow p$ for the element $\alpha_Q(p)$, and $Q \rightarrow P$, for the set $\alpha_Q(P)$, i.e., $\{Q \rightarrow p : p \in P\}$. If Q is empty, then $Q \rightarrow p = p$, and if P is empty, then $Q \rightarrow P$ also is the empty set. At last, $\{q\} \rightarrow \{p\} = \{q \rightarrow p\} = q \rightarrow p$. Observe that, if $P = \{p_1, p_2, \dots, p_m\}$, then

$$\alpha_{(Q \rightarrow P)} = \alpha_{(Q \rightarrow p_m)} \cdots \alpha_{(Q \rightarrow p_2)} \alpha_{(Q \rightarrow p_1)}. \quad (2)$$

For example, $P \rightarrow P = 1$, and if $P = \{p_1, p_2\}$ and $Q := \{q_1, q_2\}$, then

$$\begin{aligned} \alpha_{(Q \rightarrow P)} x &= \alpha_{(Q \rightarrow p_1)} \alpha_{(Q \rightarrow p_2)} x = (Q \rightarrow p_1) \rightarrow ((Q \rightarrow p_2) \rightarrow x) \\ &= (q_1 \rightarrow (q_2 \rightarrow p_1)) \rightarrow ((q_1 \rightarrow (q_2 \rightarrow p_2)) \rightarrow x). \end{aligned}$$

For further reference, we list some properties of the operations α_P , where \leq is the natural ordering of the semilattice CE^f (recall that it is defined pointwise).

Lemma 1 *In A ,*

- (a) $\alpha_P \alpha_Q = \alpha_Q \alpha_P$,
- (b) $\alpha_Q \leq \alpha_P \alpha_Q$,
- (c) $\alpha_P \leq \alpha_Q$ if and only if $\alpha_P \alpha_Q = \alpha_Q$.
- (d) $\alpha_P \alpha_Q = \alpha_{(P \cup Q)}$,
- (e) if $P \subseteq Q$, then $\alpha_P \leq \alpha_Q$,

- (f) $\alpha_Q \alpha_{(Q \rightarrow P)} = \alpha_P \alpha_Q$,
- (g) $\alpha_P \leq \alpha_Q$ iff $Q \rightarrow P = 1$.

Proof Items (a), (b) and (c) are obvious, and (e) follows from (c) and (d). In virtue of (a), items (d) and (f) generalize the Hilbert algebra identities $x \rightarrow (y \rightarrow z) = y \rightarrow (x \rightarrow z)$ and $x \rightarrow (x \rightarrow y) = x \rightarrow y$, respectively. For the “only if” part of (g), observe that $\alpha_P(p) = 1$ whenever $p \in P$. At last, if $Q \rightarrow P = 1$, then (f) and (c) lead us to left-side inequality of (g). \square

3 The adjoint semilattice of A

We first extend to Hilbert algebras a result stated for implicative semilattices in [16, Proposition 3.6].

Proposition 2 *The subset CE^f is join-dense in the poset CE , i.e., every closure endomorphism is a join of a subset of CE^f . More exactly, if $\varphi \in CE$, then $\varphi = \bigvee(\alpha_p : p \in K_\varphi)$, where K_φ is the kernel of φ .*

Proof At first, $\alpha_p \leq \varphi$ for every $p \in K_\varphi$: if $\varphi p = 1$, then, for every x , $(p \rightarrow x) \rightarrow \varphi x = \varphi((p \rightarrow x) \rightarrow x) = (\varphi p \rightarrow \varphi x) \rightarrow \varphi x = 1$ (see (1)). At second, if ψ is another upper bound of $\{\alpha_p : p \in K_\varphi\}$ and $p = \varphi a \rightarrow a$ for some $a \in A$, then $\psi a \geq \alpha_p a = p \rightarrow a \geq \varphi a$. Thus, φ is the least upper bound of the set. \square

A well-known description of the filter $[X]$ generated by some subset X of A , which goes back to [5, 14], may be formulated in terms of closure endomorphisms as follows: $a \in [X]$ if and only if a belongs to the kernel of some α_P with $P \subseteq_{\text{fin}} X$. If X is finite, one may put $P = X$. Therefore, the kernel of α_P is the filter $[P]$ generated by P ; this correspondence between endomorphisms from CE^f and finitely generated filters is bijective. By Lemma 1(g), it is even order-preserving: $Q \rightarrow P = 1$ iff $P \subseteq K_{\alpha_Q}$ iff $[P] \subseteq [Q]$. Moreover, the kernel of $\alpha_P \alpha_Q$ is the standard join of filters $[P]$ and $[Q]$, i.e., the least filter including both $[P]$ and $[Q]$. Indeed, $K_{\alpha_P}, K_{\alpha_Q} \subseteq K_{\alpha_P \alpha_Q}$ (Lemma 1(b)). Suppose that, on the other hand, $K_{\alpha_P}, K_{\alpha_Q} \subseteq K_{\alpha_R}$ for some $R \subseteq_{\text{fin}} A$. If now $x \in K_{\alpha_P \alpha_Q}$, then $\alpha_Q x \in K_{\alpha_P}$ and, further, $\alpha_Q x \in K_{\alpha_R}$. Then $\alpha_R x \in K_{\alpha_Q}$ by Lemma 1(a), and, further $\alpha_R x \in K_{\alpha_R}$, i.e., $x \in K_{\alpha_R}$ (Lemma 1(d)). Therefore, $K_{\alpha_P \alpha_Q}$ is the least upper bound of K_{α_P} and K_{α_Q} .

These considerations are summed up in the next proposition.

Proposition 3 *The transformation $\alpha_P \mapsto [P]$ is an isomorphism of CE^f onto the semilattice of finitely generated filters.*

A *subtractive semilattice* [4] is the order dual of an implicative semilattice. We are going to show that the adjoint semilattice of a Hilbert algebra A is subtractive, i.e., that there is a binary operation $-$ (*subtraction*) on CE^f such that, for all $P, Q, R \subseteq_{\text{fin}} A$,

$$\alpha_P - \alpha_Q \leq \alpha_R \text{ if and only if } \alpha_P \leq \alpha_Q \circ \alpha_R. \quad (3)$$

Taking into account Proposition 3, this fact could be derived from Theorem 2.3 of [3], where the set of finitely generated filters of a Hilbert algebra was shown to be a subtractive semilattice. The theorem itself was proved referring to several constructions from Section 6 in [7]. We shall give for the adjoint semilattice a concise direct proof.

Theorem 4 *The operation – defined on CE^f by*

$$\alpha_P - \alpha_Q := \alpha_{(Q \rightarrow P)}$$

is a subtraction.

Proof We have to prove that

$$\alpha_{(Q \rightarrow P)}x \leq \alpha_Rx \text{ for all } x \text{ if and only if } \alpha_Px \leq \alpha_Q\alpha_Rx \text{ for all } x.$$

The “only if” part holds by virtue of Lemma 1(f):

$$\alpha_P \leq \alpha_Q\alpha_P = \alpha_Q\alpha_{(Q \rightarrow P)} \leq \alpha_Q\alpha_R.$$

Conversely, from the right-side inequality, $1 = \alpha_Pp \leq \alpha_Q\alpha_Rp$ for any $p \in P$. Now observe that $\alpha_Qp \leq (\alpha_Qp \rightarrow x) \rightarrow x = \alpha_{(Q \rightarrow p)}x \rightarrow x$. Hence, for every x ,

$$1 = \alpha_R\alpha_Qp \leq \alpha_R(\alpha_{(Q \rightarrow p)}x \rightarrow x) = \alpha_{(Q \rightarrow p)}x \rightarrow \alpha_Rx$$

(see (1)) and, further, $\alpha_{(Q \rightarrow p)}x \leq \alpha_Rx$. By (2), then $\alpha_{(Q \rightarrow P)}x \leq \alpha_Rx$ for all x . \square

We next show that the transfer from Hilbert algebras to their adjoint (subtractive) semilattices is functorial. Suppose that A and A' are Hilbert algebras and that CE^f and CE'^f are the respective adjoint semilattices. Given a homomorphism $f: A \rightarrow A'$, let $f^*: CE^f \rightarrow CE'^f$ be the mapping defined by $f^*(\alpha_P) := \alpha_{f(P)}$.

Theorem 5 *Suppose that A , A' and A'' are Hilbert algebras, ε is the identity endomorphism of A , and $f: A \rightarrow A'$, $g: A' \rightarrow A''$ are homomorphisms. Then*

- (a) f^* and g^* are subtractive homomorphisms.
- (b) ε^* is the identity morphism of CE^f ,
- (c) $(gf)^* = g^*f^*$.

Proof (a) f^* is a semilattice homomorphism:

$$f^*(\alpha_P\alpha_Q) = f^*(\alpha_{P \cup Q}) = \alpha_{f(P \cup Q)} = \alpha_{f(P) \cup f(Q)} = \alpha_{f(P)}\alpha_{f(Q)} = f^*(\alpha_P)f^*(\alpha_Q),$$

and preserves subtraction: for $P = \{p_1, p_2, \dots, p_n\}$,

$$\begin{aligned} f^*(\alpha_P - \alpha_Q) &= f^*(\alpha_{(Q \rightarrow P)}) = \alpha_{f(Q \rightarrow P)} \\ &= \alpha_{(f(Q) \rightarrow f(P))} = \alpha_{f(P)} - \alpha_{f(Q)} = f^*(\alpha_P) - f^*(\alpha_Q). \end{aligned}$$

(b) is evident, as $\varepsilon = \alpha_1$.

(c) $(gf)^*(\alpha_P) = \alpha_{g(f(P))} = g^*(\alpha_{f(P)}) = g^*(f^*(\alpha_P))$. \square

Finally, we characterise the lattice of ideals of CE^f . The subsequent theorem is partly suggested by various general results in [13] on ideals of a BCI-algebra.

Theorem 6 *The filter lattice of a Hilbert algebra is isomorphic to the ideal lattice of its adjoint semilattice.*

Proof It consists of several steps. Suppose that A is a Hilbert algebra, and CE^f , its adjoint semilattice. Let I stand for the ideal lattice of the semilattice CE^f , and F , for the filter lattice of A .

(a) For every filter J of A , the subset $i(J) := \{\alpha_P : P \subseteq_{\text{fin}} J\}$ is an ideal of CE^f :

(a1) the identity closure endomorphism belongs to $i(J)$, for $\varepsilon = \alpha_1$;

(a2) if $\alpha_P, \alpha_Q \in i(J)$ with $P, Q \subseteq_{\text{fin}} J$, then also $P \cup Q \subseteq_{\text{fin}} J$ and, further, $\alpha_{P \cup Q} \in i(J)$, i.e., $\alpha_P \alpha_Q \in i(J)$;

(a3) if $\alpha_Q \in i(J)$ with $Q \subseteq_{\text{fin}} J$, and if $\alpha_P \leq \alpha_Q$ for some finite P , then $\alpha_Q x \geq \alpha_P x = 1$ and $x \in J$ for every $x \in P$. Therefore, $P \subseteq_{\text{fin}} J$ and $\alpha_P \in i(J)$.

(b) The transformation $i : F \rightarrow I$ is order-preserving: if $J \subseteq J'$, then every finite subset of J is also a subset of J' , and then $i(J) \subseteq i(J')$.

(c) For every ideal $N \in I$, the subset $j(N) := \{p : \alpha_p \in N\}$ is a filter of A :

(c1) $1 \in j(N)$, for $\alpha_1 = \varepsilon \in N$;

(c2) if p and q are compatible elements of $j(N)$ and $r := p \wedge q$, then $\alpha_p, \alpha_q \in N$, $\alpha_r = \alpha_p \circ \alpha_q \in N$ and, further, $r \in j(N)$;

(c3) if $p \in j(N)$ and $q \geq p$, then $\alpha_p \in N$, $\alpha_q \leq \alpha_p$ and, furthermore, $\alpha_q \in N$, i.e., $q \in J$.

(d) The transformation $j : I \rightarrow F$ is order-preserving: if $N \subseteq N'$, i.e., $\alpha_p \in N'$ whenever $\alpha_p \in N$, then $p \in j(N')$ for all $p \in j(N)$, and $j(N) \subseteq j(N')$.

(e) The transformations i and j are mutually inverse:

(e1) $j(i(J)) = J$: if $q \in J$, then $\alpha_q \in i(J)$ and $q \in j(i(J))$, and if $q \in j(i(J))$, then $\alpha_q \in i(J)$, i.e., $\alpha_q = \alpha_P$ for some $P \subseteq_{\text{fin}} J$. Hence, $q \in P$ and $q \in J$;

(e2) $i(j(N)) = N$: if $\alpha_P \in i(j(N))$ with $P \subseteq_{\text{fin}} j(N)$, then $\alpha_p \in N$ for all $p \in P$, and α_P , being the join of all these α_p , also belongs to N . Conversely, if $\alpha_P \in N$, then $\alpha_p \leq \alpha_P$, $\alpha_p \in N$ and $p \in j(N)$ for all $p \in P$, i.e., $P \subseteq j(N)$ and, further, $\alpha_P \in i(j(N))$.

Eventually, i and j are order isomorphisms from F to I and from I to F , respectively. Therefore, the lattices F and I are isomorphic. \square

4 Principal closure endomorphisms

Each principal filter $[p]$ is the kernel of α_p and conversely; for this reason we call closure endomorphisms α_p *principal*. Let CE^α stand for the set of all such endomorphisms. We now can say more about the transformation $p \mapsto \alpha_p$.

Theorem 7 *In A ,*

- (a) $p \leq q$ if and only if $\alpha_q \leq \alpha_p$,
- (b) $\alpha_{p \rightarrow q} = \alpha_q - \alpha_p$,
- (c) $\alpha_{p \vee q} = \alpha_p \wedge \alpha_q$ whenever $p \vee q$ exists,

- (d) $p \mathcal{C} q$ if and only $\alpha_p \circ \alpha_q$ is a principal closure endomorphism, and then $\alpha_{p \wedge q} = \alpha_p \circ \alpha_q$.
- (e) a finite subset P of A is compatible if and only if the closure endomorphism α_P is principal, and then $\alpha_{(\bigwedge P)} = \alpha_P$.

Proof (a) Clearly, if $p \leq q$, then $\alpha_q \leq \alpha_p$. If, conversely, $q \rightarrow x \leq p \rightarrow x$ for all x , then substitution of q for x shows that $p \leq q$.

(b) By the definition of subtraction.

(c) is an easy consequence of (a). Suppose that $p \vee q$ exists in A , then $\alpha_{p \vee q} \leq \alpha_p, \alpha_q$. On the other hand, if $\alpha_r \leq \alpha_p, \alpha_q$ for some r , then $p, q \leq r$ and $p \vee q \leq r$. Hence, $\alpha_r \leq \alpha_{p \vee q}$, i.e., $\alpha_{p \vee q}$ is indeed the least upper bound of α_p and α_q .

(d) If $p \wedge q$ exists in A , then similarly, $\alpha_{p \wedge q} = \alpha_p \alpha_q$. Conversely, if $\alpha_p \alpha_q = \alpha_r$ for some r , then $r \rightarrow x = p \rightarrow (q \rightarrow x)$ for all x , whence $r \leq p, q$ and $p \leq q \rightarrow r$ (put $x := p, q, r$). Thus, $r = p \wedge q$, and $p \mathcal{C} q$.

(e) If $\bigwedge P$ exists in A , then $[P] = [\bigwedge P]$ and, further $\alpha_P = \alpha_{(\bigwedge P)}$. If $\alpha_P = \alpha_r$ for some r , then $[P] = [r]$ and r is the compatible meet of P . \square

The item (d) of the theorem is, in fact, contained in Theorem 3 of [14]. In virtue of items (b) and (a), the set of principal closure endomorphisms of A turns out to be closed under subtraction and is actually an order-dual copy of A ; we shall call it the *dual algebra* of A . Therefore, CE^α is an implicative model or, as we prefer to say, a Henkin algebra. (It is now known well that the class of Henkin algebras coincides with that of positive implicative BCK-algebras described in [10].)

Corollary 8 *The set of principal closure endomorphisms of a Hilbert algebra A is a Henkin algebra dual to A .*

If every pair of elements of A is compatible, then, according to item (e) of the above theorem, all finitely generated closure endomorphisms are principal. We thus come to the following conclusion.

Corollary 9 *The adjoint semilattice of an implicative semilattice A is dually isomorphic to A .*

It follows that every subtractive semilattice is isomorphic to the adjoint semilattice of a Hilbert algebra. Also, non-isomorphic Hilbert algebras may have isomorphic adjoint semilattices. We obtain one more conclusion by help of Theorem 7(c).

Corollary 10 *If a Hilbert algebra A is an upper semilattice, then its adjoint semilattice is a sublattice of CE .*

Proof It suffices to prove that CE^f is closed under meets whenever all joins $p \vee q$ exist in A . As the lattice CE is distributive (see [2, Corollary 3.5]), for all

finite P and Q ,

$$\begin{aligned}\alpha_P \wedge \alpha_Q &= \bigvee(\alpha_p : p \in P) \wedge \bigvee(\alpha_q : q \in Q) \\ &= \bigvee(\alpha_p \wedge \alpha_q : p \in P, q \in Q) = \bigvee(\alpha_{p \vee q} : p \in P, q \in Q) = \alpha_{(P \vee Q)},\end{aligned}$$

where $P \vee Q := \{p \vee q : p \in P, q \in Q\}$. \square

The items (c) and (e) of Theorem 7 can be extended to joins and certain meets of arbitrary subsets of A . We call a subset $Y \subseteq A$ *K-compatible* if it has a lower bound which belongs to $[P]$ for every finite P such that $Y \subseteq [P]$. Let z be such a lower bound. If u is any other lower bound of Y , then $Y \subseteq [u]$ and, further, $z \in [u]$, i.e., $u \leq z$. Thus, z is the greatest lower bound; we say that it is a *K-compatible meet* of Y and denote it by $\bigwedge Y$. Evidently, if Y is finite, then this version of compatibility agrees with that discussed in Section 2: Y is K-compatible if and only if the filter generated by Y is principal, and $p = \bigwedge Y$ if and only if $[p] = [Y]$. Observe that even infinite subset Y is K-compatible, if it generates a principal filter; the converse need not hold.

Theorem 11 *Let Y be an arbitrary subset of A . Then*

- (a) $\alpha_{(\bigvee Y)} = \bigwedge\{\alpha_y : y \in Y\}$ whenever $\bigvee Y$ exists,
- (b) Y is K-compatible if and only if the set $\{\alpha_y : y \in Y\}$ has a join in CE^f that is a principal closure endomorphism, say, α_r , and then $\alpha_{(\bigwedge Y)} = \alpha_r$.

Proof (a) Similarly to item (c) of the previous theorem.

(b) By Lemma 1(g), $\alpha_q \leq \alpha_P$ iff $P \rightarrow q = 1$ iff $q \in K_{\alpha_P} = [P]$. Now suppose that r is a K-compatible meet of Y . Then $r \leq y$ for all $y \in Y$ and, for every finite P with $Y \subseteq [P]$, also $r \in [P]$. Consequently, $\alpha_y \leq \alpha_r$ for these y , i.e., α_r is an upper bound of $\{\alpha_y : y \in Y\}$. It is actually a least upper bound: if $\alpha_y \leq \alpha_P$ for all $y \in Y$ and some finite P , then $y \in [P]$; so $Y \subseteq [P]$ and, by the choice of r , $r \in [P]$, i.e., $[r] \subseteq [P]$. Now Proposition 3 implies that $\alpha_r \leq \alpha_P$.

Conversely, suppose that α_r is the join of $\{\alpha_y : y \in Y\}$. Then, in particular, $\alpha_y \leq \alpha_r$ and, by Theorem 7(a), $r \leq y$ for all $y \in Y$. Thus r is a lower bound of Y . On the other hand, if $Y \subset [P]$ for some finite P , then, for every $y \in Y$, $[y] \subseteq [P]$ and $\alpha_y \leq \alpha_P$ (Proposition 3). By choice of r , also $\alpha_r \leq \alpha_P$ and further $r \in P$. Hence, Y is K-compatible. \square

A significant consequence of this theorem will be obtained in Section 5 (Corollary 15).

5 Minimal Brouwerian extensions of A

Corollary 8 and the observations preceding it motivate a transfer from the adjoint semilattice of a Hilbert algebra to certain extensions of the latter.

We say that an implicative semilattice B is a *Brouwerian extension* of a Hilbert algebra A if A is a subreduct of B . If this is the case, then, for all $x, y, z \in A$, $z = x \wedge y$ if and only if $z = x \wedge y$ in B [3, Lemma 3.3]. Such an

extension of A is said to be *minimal*, if it is generated by A . Equivalently, B is a minimal Brouwerian extension if every its element can be presented as a join of a finite number of elements of A . Indeed, the set of those elements of B which can be so presented is closed under \rightarrow , as the following identity (with $P = \{p_1, p_2, \dots, p_m\}$) shows:

$$\bigwedge Q \rightarrow \bigwedge P = (Q \rightarrow p_1) \wedge (Q \rightarrow p_2) \wedge \dots \wedge (Q \rightarrow p_m).$$

Theorem 12 *A Brouwerian extension of a Hilbert algebra A is minimal if and only it is dually isomorphic to CE^f .*

Proof The condition is sufficient by the definition of CE^f (and Corollary 8). Now suppose that B is a minimal extension of A . Then every closure endomorphism in the adjoint semilattice of B can be presented in the form α_P with $P \subseteq_{\text{fin}} A$. Indeed, all closure endomorphisms of an implicative semilattice are principal. As any element $p \in B$ is a meet of a finite subset P of A , it follows that $\alpha_p = \alpha_P$. Now, the restriction $\alpha_P|A$ coincides with the closure endomorphism α_P of A ; thus, there is a bijection between adjoint semilattices of B and A . Furthermore, restriction to A preserves composition and subtraction and respects the identity endomorphism of B . Therefore, the adjoint semilattices are isomorphic. Hence, B is dually isomorphic to the adjoint semilattice of A . \square

Corollary 13 *Every Hilbert algebra has a unique (up to isomorphism) minimal Brouwerian extension.*

Remark 14 A construction of a minimal Brouwerian extension is implicit in A. Horn's paper [7]; see Theorem 8.5 therein (his C-algebras are just Hilbert algebras). Starting from a Hilbert algebra A , the author builds up an algebra B of non-empty subsets of A with operations \cup and \rightarrow and a constant 1 (cf. Section 2 above). The relation eq on B defined by

$$P \text{ eq } Q \quad \text{iff} \quad P \rightarrow Q = Q \rightarrow P = 1.$$

is shown to be a congruence, and the quotient algebra B/eq is an implicative semilattice. Moreover, $\{p\} \text{ eq } \{q\}$ iff $p = q$. In this way, A is embedded into an implicative semilattice. The author does not prove that the obtained extension of A is minimal; this follows from our Lemma 1(g).

Observe that $P \text{ eq } Q$ iff $\alpha_P(Q) = \alpha_Q(P) = 1$ iff $[P] = [Q]$ iff $\alpha_P = \alpha_Q$.

Due to the above theorem, we may infer several properties of minimal Brouwerian extensions from the results of previous sections. Thus, any minimal Brouwerian extension of a Hilbert algebra is an example of the implicative semilattice mentioned in the next corollary.

Corollary 15 *Every Hilbert algebra can be embedded into an implicative semilattice with preservation of arbitrary existing joins and exactly K -compatible meets.*

Remark 16 Up to order duality, this corollary is a concise version of the extension theorem for L. Henkin's implicative models which was announced by Carol R. Karp in 1954 [11] (see also the first chapter in her Ph.D. thesis [12]). The above condition of K-compatibility is just a conjunction of her conditions (i) (borrowed from [6]) and (ii), while the subcondition (2) of (i) is actually a particular case of (ii). She also observed that every implicative model is isomorphic to a subspace of closed sets of a topological space, thus anticipating the topological representation theorem of Hilbert algebras stated by A. Diego in [5].

The subsequent counterpart of [7, Theorem 8.4] is the dual of Corollary 8.

Corollary 17 *A minimal Brouwerian extension of an upper Hilbert semilattice is an implicative lattice.*

It follows from Corollary 2.4 in [3] that the filter lattice of a Hilbert algebra is isomorphic to the filter lattice of some implicative semilattice. Theorem 6 above allows us to improve this observation.

Corollary 18 *The filter lattices of a Hilbert algebra and its minimal Brouwerian extension are isomorphic.*

Namely, if A is a Hilbert algebra and B is its minimal Brouwerian extension, then, as analysis of the transformations i and j in the proof of Theorem 6 shows, the filter J^* of B which corresponds to a filter J of A is given by $J^* := \{\bigwedge P : P \subseteq_{\text{fin}} J\}$, and then $J = J^* \cap A$.

References

- [1] Cīrulis, J.: *Multipliers in implicative algebras*. Bull. Sect. Log. (Łódź) **15** (1986), 152–158.
- [2] Cīrulis, J.: Multipliers, closure endomorphisms and quasi-decompositions of a Hilbert algebra. In: Chajda et al. I. (eds) *Contrib. Gen. Algebra Verlag Johannes Heyn*, Klagenfurt, 2005, 25–34.
- [3] Cīrulis, J.: *Hilbert algebras as implicative partial semilattices*. Centr. Eur. J. Math. **5** (2007), 264–279.
- [4] Curry, H. B.: Foundations of Mathematical logic. *McGraw-Hill*, New York, 1963.
- [5] Diego, A.: Sur les algèbres de Hilbert. *Gauthier-Villars; Nauwelaerts*, Paris; Louvain, 1966.
- [6] Henkin, L.: *An algebraic characterization of quantifiers*. Fund. Math. **37** (1950), 63–74.
- [7] Horn, A.: *The separation theorem of intuitionistic propositional calculus*. Journ. Symb. Logic **27** (1962), 391–399.
- [8] Huang, W., Liu, F.: *On the adjoint semigroups of p-separable BCI-algebras*. Semigroup Forum **58** (1999), 317–322.
- [9] Huang, W., Wang, D.: *Adjoint semigroups of BCI-algebras*. Southeast Asian Bull. Math. **19** (1995), 95–98.
- [10] Iseki, K., Tanaka, S.: *An introduction in the theory of BCK-algebras*. Math. Japon. **23** (1978), 1–26.

- [11] Karp, C. R.: *Set representation theorems in implicative models*. Amer. Math. Monthly **61** (1954), 523–523 (abstract).
- [12] Karp, C. R.: Languages with expressions of infinite length. *Univ. South. California*, 1964 (Ph.D. thesis).
- [13] Kondo, M.: *Relationship between ideals of BCI-algebras and order ideals of its adjoint semigroup*. Int. J. Math. **28** (2001), 535–543.
- [14] Marsden, E. L.: *Compatible elements in implicational models*. J. Philos. Log. **1** (1972), 195–200.
- [15] Schmidt, J.: *Quasi-decompositions, exact sequences, and triple sums of semigroups I. General theory. II Applications*. In: Contrib. Universal Algebra Colloq. Math. Soc. Janos Bolyai (Szeged) **17** North-Holland, Amsterdam, 1977, 365–428.
- [16] Tsinakis, C.: *Brouwerian semilattices determined by their endomorphism semigroups*. Houston J. Math. **5** (1979), 427–436.
- [17] Tsirulish, Ya. P.: *Notes on closure endomorphisms of implicative semilattices*. Latvijiskij Mat. Ezhegodnik **30** (1986), 136–149 (in Russian).