

GLOBAL LIMIT CYCLE BIFURCATIONS OF LOW-DIMENSIONAL DYNAMICAL SYSTEMS

VALERY GAIKO

United Institute of Informatics Problems, National Academy of Sciences of Belarus

Leonid Beda Str. 6-4, Minsk 220040, Belarus

E-mail: valery.gaiko@gmail.com

The qualitative analysis of polynomial dynamical systems is carried out. To control all of their limit cycle bifurcations, especially, bifurcations of multiple limit cycles, it is necessary to know the properties and combine the effects of all of their rotation parameters. It can be done by means of the development of new bifurcational and topological methods based on the well-known Weierstrass preparation theorem and the Perko planar termination principle stating that the maximal one-parameter family of multiple limit cycles terminates either at a singular point which is typically of the same multiplicity (cyclicity) or on a separatrix cycle which is also typically of the same multiplicity (cyclicity) [1]. This principle is a consequence of the principle of natural termination which was stated for higher-dimensional dynamical systems by A. Wintner who studied one-parameter families of periodic orbits of the restricted three-body problem and used Puiseux series to show that in the analytic case any one-parameter family of periodic orbits can be uniquely continued through any bifurcation except a period-doubling bifurcation. Such a bifurcation can happen, e. g., in a three-dimensional Lorenz system. This cannot happen for planar systems. That is why the Wintner–Perko termination principle is applied for studying multiple limit cycle bifurcations of planar polynomial dynamical systems [1].

If we do not know the cyclicity of the termination points, then, applying canonical systems with field rotation parameters, we use geometric properties of the spirals filling the interior and exterior domains of limit cycles. Applying this method, we have solved, e. g., Smale’s Thirteenth Problem proving that the Liénard system with a polynomial of degree $2k + 1$ can have at most k limit cycles. Generalizing the obtained results, we have also solved the problem of the maximum number of limit cycles surrounding a singular point for an arbitrary polynomial system and Hilbert’s Sixteenth Problem for a general Liénard polynomial system with an arbitrary (but finite) number of singular points [2]. Applying a similar approach, we complete the strange attractor bifurcation scenario which connects globally the homoclinic, period-doubling, Andronov–Shilnikov, and period-halving bifurcations in the classical Lorenz system [3]. We discuss also how to apply this approach for studying global limit cycle bifurcations of discrete polynomial (and rational) dynamical systems which model the population dynamics in biomedical and ecological systems.

This work was partially supported by the Simons Foundation of the International Mathematical Union and the Department of Mathematics and Statistics of the Missouri University of Science and Technology.

REFERENCES

- [1] V. A. Gaiko. *Global Bifurcation Theory and Hilbert’s Sixteenth Problem*. Kluwer, Boston, 2003.
- [2] V. A. Gaiko. The applied geometry of a general Liénard polynomial system. *Appl. Math. Letters*, **25** (12): 2327–2331, 2012.
- [3] V. A. Gaiko. Global bifurcation analysis of the Lorenz system. *J. Nonlinear Sci. Appl.*, **7** (6): 429–434, 2014.